


ILD/IPF SHOULD BE CONSIDERED IN ALL PATIENTS WITH CHRONIC COUGH AND EXERTIONAL DYSPNEA¹

 **OVER 50%**

OF PATIENTS WITH IPF ARE INITIALLY MISDIAGNOSED WITH OTHER COMMON RESPIRATORY DISEASES⁹

Use this tool to help assess when a diagnosis of ILD/IPF should be pursued



CHF, congestive heart failure; COPD, chronic obstructive pulmonary disease; ILD, interstitial lung disease; IPF, idiopathic pulmonary fibrosis; PFT, pulmonary function test.

References: 1. Raghu G et al; on behalf of the ATS/ERS/JRS/ALAT Committee on Idiopathic Pulmonary Fibrosis. *Am J Respir Crit Care Med.* 2011;183(6):788-824. 2. American Thoracic Society. *Am J Respir Crit Care Med.* 1995;152(5 Pt 2):S77-S121. 3. Mayo Clinic. Heart failure symptoms. <http://www.mayoclinic.org/diseases-conditions/heartfailure/basics/symptoms/con-20029801>. Accessed December 2, 2021. 4. Ryu JH et al. *Mayo Clin Proc.* 2007;82(8):976-986. 5. Niimi A. *Curr Respir Med Rev.* 2011;7(1):47-54. 6. Kaufman J. *Nurse management.* <https://booksite.elsevier.com/samplechapters/9780323065801/9780323065801.pdf>. Accessed December 2, 2021. 7. Johnson JD et al. *Am Fam Physician.* 2014;89(5):359-366. 8. von Cleemput J et al. *Adv Ther.* 2019;36(2):298-317. doi:10.1007/s12325-018-0857-z. 9. Collard HR et al. *Respir Med.* 2007;101:1350-1354.